

A CONVEXITY THEOREM FOR THREE TANGLED HAMILTONIAN TORUS ACTIONS

HIRAKU ABE

Abstract The convexity theorem for Hamiltonian torus actions states that the image of a moment map of a Hamiltonian torus action on a compact, connected symplectic manifold is a convex polytope ([A], [G-S]). Kirwan generalized this theorem to the case of any compact, connected Lie group, which also gives us a convex polytope ([K]). On the other hand, if a torus which has half the dimension of the manifold acts effectively in a Hamiltonian fashion, then its moment map provides completely integrable systems. Many important completely integrable systems are super-integrable systems in which each trajectory is contained in a smaller torus than the Liouville torus: harmonic oscillators, the Kepler system, the Toda lattice, etc. In this talk, motivated by the structure of the super-integrable system on the Toda lattice ([A-D-S]), a new generalization of the convexity theorem for Hamiltonian torus actions will be demonstrated.

REFERENCES

- [A] Atiyah, M. F. *Convexity and commuting Hamiltonians*, Bull. Lond. Math. Soc. **14** (1982), 1-15.
- [A-D-S] M. Agrotis, P. A. Damianou, and C. Sophocleous, *The Toda lattice is super-integrable*, arXiv:math-ph/0507051v1.
- [G-S] Guillemin, V. and Sternberg, S. *Convexity properties of the moment mapping*, Invent. math. **67** (1982), 491-513.
- [K] Kirwan, F. *Convexity properties of the moment mapping III*, Invent. math. **77** (1984), 547-552.