RR = #BS VIA LOCALIZATION OF INDEX

TAKAHIKO YOSHIDA

For a pre-quantized closed symplectic manifold a Riemann- $Roch\ number$ is defined to be the index of a $Spin^c$ Dirac operator with coefficients in the pre-quantization line bundle.

Suppose the pre-quantized symplectic manifold is equipped with a structure of a Lagrangian fiber bundle. A fiber of the Lagrangian fiber bundle is said to be *Bohr-Sommerfeld* if the restriction of the pre-quantization line bundle to the fiber is trivially flat. Bohr-Sommerfeld fibers appear discretely. Then Andersen showed in [1] that the Riemann-Roch number is equal to the number of Bohr-Sommerfeld fibers.

Similar phenomena have been observed for several examples of Lagrangian fiber bundles with singular fibers, such as,

- moment maps of toric varieties [2],
- the Gelfand-Cetlin system for complex flag manifolds [7],
- Goldman's Hamiltonian system on the moduli space of flat SU(2)-bundle on a Riemann surface [5, 9],

and for the following generalizations

- pre-symplectic toric manifolds [10],
- Spin^c manifolds [6],
- torus manifolds [11, 8],

and so on.

In this series of lectures I explain an approach to investigate these phenomena by using a localization of index. The plan of this lectures is as follows

- § 1 Hamiltonian mechanics on symplectic manifolds
- § 2 Lagrange fiber bundles
- § 3 Quantization
- § 4 An approach to RR= #BS by localization of index.

In \S 1 I briefly explain the Hamiltonian mechanics on symplectic manifolds. \S 1 is an introduction to symplectic manifolds and also necessary for \S 3. In \S 2 I introduce a notion of a Lagrange fiber bundle and prove its classification theorem. In \S 3 I explain some background of the above phenomena. In \S 4 I explain a localization technique of the index of a Dirac-type operator and its application to RR= #BS. \S 4 is based on a joint work [3, 4] with Mikio Furuta and Hajime Fujita. I show that if a prequantized symplectic manifold admits a structure of a singular Lagrangian fiber bundle, then, the Riemann-Roch number is localized on nonsingular Bohr-Sommerfeld fibers and singular fibers. The idea used here is the Witten deformation that is used to prove the Morse inequalities in [12].

1

References

- 1. J. E. Andersen, Geometric quantization of symplectic manifolds with respect to reducible non-negative polarizations, Comm. Math. Phys. 183 (1997), no. 2, 401–421.
- 2. V. Danilov, *The geometry of toric varieties (Russian)*, Uspekhi Mat. Nauk **33** (1978), no. 2, 85–134, English translation: Russian Math. Surveys 33 (1978), no. 2, 97–154.
- 3. H. Fujita, M. Furuta, and T. Yoshida, Torus fibrations and localization of index I, UTMS Preprint Series 2008-10, 23 pages. Also available at arXiv:0804.3258, 2008.
- 4. ______, Torus fibrations and localization of index II, UTMS Preprint Series 2009-21, 64 pages. Also available at arXiv:0910.0358, 2009.
- 5. W. M. Goldman, Invariant functions on Lie groups and Hamiltonian flows of surface group representations, Invent. Math. 85 (1986), no. 2, 263–302.
- 6. M. D. Grossberg and Y. Karshon, Equivariant index and the moment map for completely intergrable torus actions, Adv. Math. 133 (1998), no. 2, 185–223.
- 7. V. Guillemin and S. Sternberg, The Gelfand-Cetlin system and quantization of the complex flag manifolds, J. Funct. Anal. **52** (1983), no. 1, 106–128.
- 8. A. Hattori and M. Masuda, Theory of multi-fans, Osaka J. Math. 40 (2003), no. 1, 1–68.
- 9. L. Jeffrey and J. Weitsman, Bohr-Sommerfeld orbits in the moduli space of flat connections and the Verlinde dimension formula, Comm. Math. Phys. 150 (1992), no. 3, 593–630.
- Y. Karshon and S. Tolman, The moment map and line bundles over presymplectic toric manifolds, J. Differential Geom. 38 (1993), no. 3, 465–484.
- 11. M. Masuda, Unitary toric manifolds, multi-fans and equivariant index, Tohoku Math. J. (2) **51** (1999), no. 2, 237–265.
- 12. E. Witten, Supersymmetry and Morse theory, J. Diff. Geom. 17 (1982), no. 4, 197-229.

DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE AND TECHNOLOGY, MEIJI UNIVERSITY, 1-1 HIGASHIMITA 1-CHOME, TAMA-KU, KAWASAKI, 214-8571, JAPAN *E-mail address*: takahiko@math.meiji.ac.jp