

Vector bundles over normal varieties trivialized by finite morphisms

MARCO ANTEI AND VIKRAM B. MEHTA

Abstract. Let Y be a normal and projective variety over an algebraically closed field k and V a vector bundle over Y . We prove that if there exist a k -scheme X and a finite surjective morphism $g : X \rightarrow Y$ that trivializes V then V is essentially finite.

Mathematics Subject Classification (2010). Primary 14L15, 14L30;
Secondary 14J60.

Keywords. Essentially finite vector bundles, Finite morphisms,
Principal bundles.

1. Introduction. Essentially finite vector bundles over a reduced, connected and proper scheme Y over a perfect field k have been defined by Nori [5, 6]. They turn out to be those vector bundles V over Y which are trivialized by some principal G bundle $f : Z \rightarrow Y$ for a certain finite k -group scheme G (i.e., $f^*(V)$ is trivial). The aim of this paper is to prove the following

Theorem 1.1. (Cf. Theorem 2.1). *Let k be any algebraically closed field and Y a projective and normal variety over k . Assume there exist a projective variety X over k and a finite surjective morphism $g : X \rightarrow Y$ such that $g^*(V)$ is trivial, then V is essentially finite.*

When Y is smooth then Theorem 1.1 is well known: it has first been proved by Parameswaran and Subramanian [7, Section 3], for $\dim(Y) = 1$ provided g is separable. Then it has been subsequently proved by Balaji and Parameswaran [1, Section 6] for Y smooth and projective of any dimension provided g is separable. Then finally Biswas and Dos Santos [2] have given a different proof: for any finite and surjective $g : X \rightarrow Y$, with Y smooth and projective over k , they first explain how to reduce to the case of curves ([2, Section 3]) by means of the Grothendieck–Lefschetz theorem for the S -fundamental group

scheme, then in loc. cit. Section 4.2 they prove Theorem 1.1 for Y a smooth and projective curve and $g : X \rightarrow Y$ separable (the crucial point) and finally they prove the result for any g (loc. cit. Section 4.3) and Y a smooth and projective curve.

Our proof of Theorem 1.1 not only holds for Y normal but it is shorter, and we use neither Tannakian categories nor Grothendieck–Lefschetz theorem for the fundamental group scheme. The main argument of our proof is Lemma 2.5, which is of independent interest, where we prove that a finite and surjective morphism $g : X \rightarrow Y$ between normal and projective varieties over any algebraically closed field k , étale outside a closed set of codimension 2 in Y , factors through a Galois étale cover $g' : X' \rightarrow Y$ if and only if there exists a non trivial vector bundle V on Y such that $g^*(V)$ is trivial on X .

2. The theorem. Throughout the whole paper k will be an algebraically closed field and Y a normal and projective variety over k . Let us denote by $EF(Y)$ the neutral Tannakian category of essentially finite vector bundles over Y . The aim of this paper is to prove the following

Theorem 2.1. *Assume there exist a normal projective variety X over k and a finite surjective morphism $g : X \rightarrow Y$ such that $g^*(V)$ is trivial, then $V \in EF(Y)$.*

It is clear that Theorem 1.1 can be easily deduced from Theorem 2.1 simply normalizing an irreducible component of X_{red} dominating Y .

Remark 2.2. This theorem holds in both zero and positive characteristic.

As pointed out in the introduction, the crucial point in the proof of Theorem 2.1 is to prove the statement for g separable (or generically étale, i.e., the extension $K(Y) \subset K(X)$ of the function fields induced by g is separable), and this will be the object of Lemma 2.4.

So first we consider the easier case where $g : X \rightarrow Y$ is purely inseparable (i.e., the extension $K(Y) \subset K(X)$ of their function fields is purely inseparable, which only occurs when $\text{char}(k) > 0$) and then it will only remain to explain how to reduce to these two cases, the separable and purely inseparable ones.

Lemma 2.3. *Assume there exist a normal projective variety X over k and a finite, surjective, purely inseparable morphism $g : X \rightarrow Y$ such that $g^*(V)$ is trivial, then $V \in EF(Y)$.*

Proof. We are in the case $\text{char}(k) = p > 0$. So let us denote by $F_X : X \rightarrow X$ and $F_Y : Y \rightarrow Y$ respectively the absolute Frobenius morphisms of X and Y . Since $K(Y) \subset K(X)$ is purely inseparable then there exists a positive integer n such that $K(X)^{(p^n)} \subset K(Y)$. This implies that there is a morphism $h : Y \rightarrow X$ such that $gh = F_Y^n$ (i.e. the Frobenius iterated n times) and $hg = F_X^n$. By assumption $g^*(V)$ is trivial on X , thus $h^*g^*(V) = (gh)^*(V) = (F_Y^n)^*(V)$ is trivial hence V is essentially finite (cf. [4, Section 2]). \square

Lemma 2.4. *Assume there exist a normal projective variety X over k and a finite, surjective, separable morphism $g : X \rightarrow Y$ such that $g^*(V)$ is trivial, then $V \in EF(Y)$.*

Proof. We may assume that $K(X)$ is normal (then Galois) over $K(Y)$ with Galois group G (if it is not simply consider the normal closure of the extension $K(Y) \subset K(X)$).

Let $W := (g_*\mathcal{O}_X)_{\max}$ be the maximal semistable subsheaf of $g_*\mathcal{O}_X$ (i.e. the first term of the Harder–Narasimhan filtration of $g_*\mathcal{O}_X$, [1, Section 6]) then its slope $\mu(W) = \mu_{\max}(g_*\mathcal{O}_X) = 0$: indeed since there is at least the canonical morphism $\mathcal{O}_Y \rightarrow g_*\mathcal{O}_X$ then in particular we have

$$0 = \mu(\mathcal{O}_Y) \leq \mu_{\max}(g_*\mathcal{O}_X);$$


but g is separable then $g^*(W)$ is still semistable; now consider the isomorphism

$$\mathrm{Hom}_X(g^*(W), \mathcal{O}_X) \simeq \mathrm{Hom}_Y(W, g_*\mathcal{O}_X) \neq 0$$

from which we deduce $\mu(g^*(W)) \leq 0$ hence $\mu(W) \leq 0$ (recall that $\mu(W) = \mu(g^*(W))/\mathrm{deg}(g)$).

The coherent sheaf W is in general only torsion free over Y . But it is locally free if restricted to a big open subset $Y_0 \subset Y$, i.e., $\mathrm{codim}_Y(Y \setminus Y_0) \geq 2$. Let $W_0 := W|_{Y_0}$ denote the vector bundle over Y_0 , $\mathrm{Sym}^*(W_0^*)$ the symmetric algebra of the dual of W_0 , and consider $X_0 := \mathbf{Spec}(\mathrm{Sym}^*(W_0^*))$ with its canonical map $g_0 : X_0 \rightarrow Y_0$.

The vector bundle W_0 is strongly semistable of degree 0 over Y_0 : indeed let us denote by F_{X_0} and F_{Y_0} respectively the absolute Frobenius morphisms of X_0 and Y_0 , and assume W is not strongly semistable then there exists a subsheaf U of $F_{Y_0}^*(W)$ such that $\mathrm{deg}(U) > 0$. Let X_1 be the fiber product of $g_0 : X_0 \rightarrow Y_0$ and F_{Y_0} . It is an integral scheme. We denote by $\mathrm{pr} : X_1 \rightarrow X_0$ and $g_1 : X_1 \rightarrow Y_0$ the projections and also $h : X_0 \rightarrow X_1$ the map given by the universal property of the fiber product:



Now $U \subseteq F_{Y_0}^*(g_{0*}(\mathcal{O}_{X_0})) = g_{1*}(\mathrm{pr}^*(\mathcal{O}_{X_0})) = g_{1*}(\mathcal{O}_{X_1})$. But from $\mathcal{O}_{X_1} \hookrightarrow h_*(\mathcal{O}_{X_0})$ we obtain $g_{1*}(\mathcal{O}_{X_1}) \hookrightarrow g_{1*}(h_*(\mathcal{O}_{X_0})) = g_{0*}(\mathcal{O}_{X_0})$ the latter being semistable whence a contradiction. As a consequence we have a homomorphism of \mathcal{O}_{Y_0} -algebras $g_{0*}(\mathcal{O}_{X_0}) \simeq W_0$ (cf. also [1, Section 6]).

Since $g_{0*}(\mathcal{O}_{X_0})$ is semistable of slope 0 over Y_0 then X_0 is a Galois-étale cover over Y_0 ([1, Lemma 6.2]), the Galois group of g_0 still being G . Now let us fix some notations: recall that by assumption V is a vector bundle over Y such that $T := g^*(V)$ is trivial on X ; we set $V_0 := V|_{Y_0}$ and $T_0 := g_0^*(V_0)$ so the latter is also trivial on X_0 . Since g_0 is a Galois-étale cover, T_0 is a G -bundle on X_0 . But X_0 is a big open set in X thus G acts on X and then G acts also on T . Since T is a G -bundle, we go on as follows: we have $X/G \simeq Y$ and

the trivial bundle T on X descends to Y . So by Kempf’s lemma (cf. e.g., [3, Théorème 2.3]), for all x in X , the stabilizer G_x acts trivially on the fibre T_x . But T is trivial and both X and X_0 have no global sections except constants, this means that there is a map

$$\rho : G \rightarrow GL(T_x) = GL_r$$

over X , where $r := \text{rank}(T)$. Assume first that the map $\rho : G \rightarrow GL_r$ is injective. We already know that G acts freely on X_0 . So let us take $x \in X \setminus X_0$: since G_x is a subgroup of G , G_x has to be trivial. This proves that G acts freely on X . So $g : X \rightarrow Y$ is a Galois-étale cover hence V is in $EF(Y)$. Up to now we have assumed ρ to be injective. If it is not, then just consider $H := G/\ker(\rho)$ and $X' = X/(\ker(\rho))$, which is provided with a faithful H -action and clearly $Y \simeq X'/H$. Hence $H \rightarrow GL_r$ is injective, V is trivial over X' and we proceed as before. □

From the previous discussion follows Lemma 2.5 which is of independent interest:

Lemma 2.5. *Let Y and X be normal and projective varieties and $f : X \rightarrow Y$ a finite and surjective morphism, étale outside a closed set of codimension 2 in Y , then we have proved that there exists a non-trivial vector bundle V on Y such that f^*V is trivial on X if and only if f factors through a Galois étale cover $f' : X' \rightarrow Y$.*

Proof. The Galois étale cover $f' : X' \rightarrow Y$ is the one constructed in the proof of Lemma 2.4. □

Remark 2.6. Let notations be as in Lemma 2.4 and its proof. In Lemma 2.5 we obtain the smallest Galois étale cover where V becomes trivial. Indeed $f' : X' \rightarrow Y$ determines and is determined by the kernel of $\rho : G \rightarrow GL_r$; if ρ is injective then $X' = X$. If ρ is not injective then $X' := X/\ker(\rho)$ is Galois étale over Y . It can happen that there are no Galois étale covers between X and X' . This happens if and only if $\mu_{\max}C < 0$, where C is the cokernel of $f'_*(\mathcal{O}_{X'}) \rightarrow f_*(\mathcal{O}_X)$.

We are now ready to prove the main result:

Proof of Theorem 2.1. if $\text{char}(k) = 0$, then Lemma 2.4 is sufficient to conclude. So let us assume $\text{char}(k) = p > 0$: if g is purely inseparable, then Lemma 2.3 is enough to conclude. Otherwise, if g is arbitrary, we argue as follows: again we may assume that $K(X)$ is normal over $K(Y)$ with Galois group G . It is known that $L := K(X)^G$ is a proper purely inseparable field extension of $K(Y)$ while $K(X)$ is separable over L , then Galois. Let Z be the integral closure of Y in L , then $g : X \rightarrow Y$ factors through the maps $s : X \rightarrow Z$ and $t : Z \rightarrow Y$ (i.e., $ts = g$) where $t : Z \rightarrow Y$ is purely inseparable and $s : X \rightarrow Z$ is separable. By Lemma 2.4 the vector bundle $W := t^*(V) \in EF(Z)$ because $s^*(W)$ is trivial on X . As we did for Lemma 2.3, there exists a morphism $h : Y \rightarrow Z$ such that $h^*t^*(V) = (th)^*(V) = (F_Y^n)^*(V)$ for some integer n ; but $h^*(W) \in EF(Y)$ thus $(F_Y^n)^*(V) \in EF(Y)$ then there exists $m \geq n$ such that $(F_Y^m)^*(V)$ is Galois-étale trivial (i.e., there exists a Galois-étale cover $j : Y' \rightarrow Y$ such

that $j^*((F_Y^m)^*(V))$ is trivial on Y' , and that is enough to conclude that V is essentially finite on Y . \square

Acknowledgements. M. Antei would like to thank Vikram Mehta for the invitation to the Tata Institute of Fundamental Research of Mumbai where this paper has been conceived. V. B. Mehta would like to thank the ICTP, Trieste for hospitality. We have been informed that V. Balaji and A. J. Parameswaran are also independently considering the same problem.

References

- [1] V. BALAJI AND A. J. PARAMESWARAN, An Analogue of the Narasimhan-Seshadri Theorem and some Applications. *J. Topol.* **4** (2011), 105–140.
- [2] I. BISWAS AND J. P. P. DOS SANTOS, Vector Bundles Trivialized by Proper Morphisms and the Fundamental Group Scheme, *J. Inst. Math. Jussieu* **10** (2011), 225–234.
- [3] J.-M. DREZET AND M. S. NARASIMHAN, Groupe de Picard des variétés de modules de fibrés semi-stables sur les courbes algébriques, *Invent. math.* **97** (1989), 53–94.
- [4] V. B. MEHTA AND S. SUBRAMANIAN, On the fundamental group scheme, *Invent. math.* **148** (2002), 143–150.
- [5] M. V. NORI, On the Representations of the Fundamental Group, *Compositio Mathematica*, **33** (1976), 29–42.
- [6] M. V. NORI, The Fundamental Group-Scheme, *Proc. Indian Acad. Sci. (Math. Sci.)*, **91** (1982), 73–122.
- [7] A. J. PARAMESWARAN AND S. SUBRAMANIAN, On the spectrum of asymptotic slopes, in: “Teichmüller Theory and moduli problems” Ramanujam Mathematical Society Lecture Notes **10** (2010), 519–528.

MARCO ANTEI

Department of Mathematical Science,
335 Gwahangno (373-1 Guseong-dong),
Yuseong-gu,
Daejeon 305-701,
Republic of Korea
e-mail: marco.antei@gmail.com

VIKRAM B. MEHTA

School of Mathematics,
Tata Institute of Fundamental Research,
Homi Bhabha Road,
Bombay 400005,
India
e-mail: vikram@math.tifr.res.in

Received: 18 May 2011